



Statistical Learning II

Lecture 5 - Least squares (continued)

Bruno Loureiro


@ CSD, DI-ENS & CNRS

brloureiro@gmail.com

Geometrical interpretation

This gives a natural interpretation of the OLS predictor as an orthogonal projection of the labels in the row space of X:

$$\hat{\boldsymbol{\theta}}_{OLS} = X^{+}y$$
 \Rightarrow $\hat{\boldsymbol{y}}_{OLS} = X\hat{\boldsymbol{\theta}}_{OLS} = XX^{+}y$



Statistical analysis of OLS

Fixed-design analysis

We now assume the following data generative model:

$$y_i = \langle \boldsymbol{\theta}_{\star}, \boldsymbol{x}_i \rangle + \varepsilon_i$$

With: • Fixed $\theta_{\star} \in \mathbb{R}^d$ and $x_i \in \mathbb{R}^d$ "fixed design"

• $\mathbb{E}[\varepsilon_i] = 0$ and $\mathbb{E}[\varepsilon_i^2] = \sigma^2 < \infty$

We now assume the following data generative model:

$$y_i = \langle \boldsymbol{\theta}_{\star}, \boldsymbol{x}_i \rangle + \varepsilon_i$$

With: • Fixed $\theta_{\star} \in \mathbb{R}^d$ and $x_i \in \mathbb{R}^d$ "fixed design"

• $\mathbb{E}[\varepsilon_i] = 0$ and $\mathbb{E}[\varepsilon_i^2] = \sigma^2 < \infty$

Remarks: • The Bayes predictor and error are given by

$$f_{\star}(x) = \mathbb{E}[y | X = x] = \langle \theta_{\star}, x \rangle$$
 $\mathcal{R}_{\star} = \mathcal{R}(\theta_{\star}) = \sigma^2$

We now assume the following data generative model:

$$y_i = \langle \boldsymbol{\theta}_{\star}, \boldsymbol{x}_i \rangle + \varepsilon_i$$

With: • Fixed $\theta_{\star} \in \mathbb{R}^d$ and $x_i \in \mathbb{R}^d$ "fixed design"

•
$$\mathbb{E}[\varepsilon_i] = 0$$
 and $\mathbb{E}[\varepsilon_i^2] = \sigma^2 < \infty$

<u>Remarks:</u> • The Bayes predictor and error are given by

$$f_{\star}(x) = \mathbb{E}[y | X = x] = \langle \theta_{\star}, x \rangle$$
 $\mathcal{R}_{\star} = \mathcal{R}(\theta_{\star}) = \sigma^2$

In particular

$$f_{\star} \in \mathcal{H} = \{ f(x) = \langle \boldsymbol{\theta}, x \rangle : \boldsymbol{\theta} \in \mathbb{R}^d \}$$


"Well-specified setting"

We now assume the following data generative model:

$$y_i = \langle \boldsymbol{\theta}_{\star}, \boldsymbol{x}_i \rangle + \varepsilon_i$$

With: • Fixed $\theta_{\star} \in \mathbb{R}^d$ and $x_i \in \mathbb{R}^d$ "fixed design"

• $\mathbb{E}[\varepsilon_i] = 0$ and $\mathbb{E}[\varepsilon_i^2] = \sigma^2 < \infty$




We now assume the following data generative model:

$$y_i = \langle \boldsymbol{\theta}_{\star}, \boldsymbol{x}_i \rangle + \varepsilon_i$$

With: • Fixed $\theta_{\star} \in \mathbb{R}^d$ and $x_i \in \mathbb{R}^d$ "fixed design"

• $\mathbb{E}[\varepsilon_i] = 0$ and $\mathbb{E}[\varepsilon_i^2] = \sigma^2 < \infty$



Given a batch of data sampled from this model:

$$y = X\theta_{\star} + \varepsilon \in \mathbb{R}^n$$

$$\hat{\boldsymbol{\theta}}_{OLS}(\boldsymbol{X}, \boldsymbol{y}) = (\boldsymbol{X}^{\mathsf{T}} \boldsymbol{X})^{-1} \boldsymbol{X}^{\mathsf{T}} \boldsymbol{y}$$

Given a batch of data sampled from this model:

$$y = X\theta_{\star} + \varepsilon \in \mathbb{R}^n$$

$$\hat{\boldsymbol{\theta}}_{OLS}(\boldsymbol{X}, \boldsymbol{y}) = (\boldsymbol{X}^{\mathsf{T}} \boldsymbol{X})^{-1} \boldsymbol{X}^{\mathsf{T}} \boldsymbol{y}$$

Given a batch of data sampled from this model:

$$y = X\theta_{\star} + \varepsilon \in \mathbb{R}^n$$

$$\hat{\boldsymbol{\theta}}_{OLS}(\boldsymbol{X}, \boldsymbol{y}) = (\boldsymbol{X}^{\top} \boldsymbol{X})^{-1} \boldsymbol{X}^{\top} \boldsymbol{y} = (\boldsymbol{X}^{\top} \boldsymbol{X})^{-1} \boldsymbol{X}^{\top} (\boldsymbol{X} \boldsymbol{\theta}_{\star} + \boldsymbol{\varepsilon})$$

Given a batch of data sampled from this model:

$$y = X\theta_{\star} + \varepsilon \in \mathbb{R}^n$$

$$\hat{\boldsymbol{\theta}}_{OLS}(X, y) = (X^{\top}X)^{-1}X^{\top}y = (X^{\top}X)^{-1}X^{\top}(X\boldsymbol{\theta}_{\star} + \boldsymbol{\varepsilon})$$
$$= (X^{\top}X)^{-1}X^{\top}X\boldsymbol{\theta}_{\star} + (X^{\top}X)^{-1}X^{\top}\boldsymbol{\varepsilon}$$

Given a batch of data sampled from this model:

$$y = X\theta_{\star} + \varepsilon \in \mathbb{R}^n$$

Our goal is to understand the statistical properties of OLS. For simplicity, assume that rank(X) = d (n > d):

$$\hat{\boldsymbol{\theta}}_{OLS}(X, y) = (X^{\top}X)^{-1}X^{\top}y = (X^{\top}X)^{-1}X^{\top}(X\boldsymbol{\theta}_{\star} + \boldsymbol{\varepsilon})$$

$$= (X^{\top}X)^{-1}X^{\top}X\boldsymbol{\theta}_{\star} + (X^{\top}X)^{-1}X^{\top}\boldsymbol{\varepsilon}$$

$$= \boldsymbol{\theta}_{\star} + \frac{1}{n}\hat{\boldsymbol{\Sigma}}_{n}^{-1}X^{\top}\boldsymbol{\varepsilon}$$

Where we have defined $\hat{\Sigma}_n = \frac{1}{n} X^{\top} X \in \mathbb{R}^{d \times d}$ (Empirical covariance)

$$\hat{\boldsymbol{\theta}}_{OLS}(\boldsymbol{X},\boldsymbol{y}) = \boldsymbol{\theta}_{\star} + \frac{1}{n}\hat{\boldsymbol{\Sigma}}_{n}^{-1}\boldsymbol{X}^{\mathsf{T}}\boldsymbol{\varepsilon}$$
"signal" "noise"

$$\hat{\boldsymbol{\theta}}_{OLS}(\boldsymbol{X},\boldsymbol{y}) = \boldsymbol{\theta}_{\star} + \frac{1}{n}\hat{\boldsymbol{\Sigma}}_{n}^{-1}\boldsymbol{X}^{\mathsf{T}}\boldsymbol{\varepsilon}$$
"signal" "noise"

In particular:

• Bias:

$$\mathbb{E}_{\varepsilon} \left[\hat{\boldsymbol{\theta}}_{OLS}(\boldsymbol{X}, \boldsymbol{y}) \right] = \boldsymbol{\theta}_{\star}$$

"Unbiased"

$$\hat{\boldsymbol{\theta}}_{OLS}(\boldsymbol{X},\boldsymbol{y}) = \boldsymbol{\theta}_{\star} + \frac{1}{n}\hat{\boldsymbol{\Sigma}}_{n}^{-1}\boldsymbol{X}^{\mathsf{T}}\boldsymbol{\varepsilon}$$
"signal" "noise"

In particular:

• Bias:
$$\mathbb{E}_{\boldsymbol{\varepsilon}}\left[\hat{\boldsymbol{\theta}}_{OLS}\!(\boldsymbol{X},\boldsymbol{y})\right] = \boldsymbol{\theta}_{\star}$$
 "Unbiased"

• Variance:
$$\operatorname{Var}_{\boldsymbol{\varepsilon}} \left[\hat{\boldsymbol{\theta}}_{OLS}(\boldsymbol{X}, \boldsymbol{y}) \right] = \frac{\sigma^2}{n} \hat{\boldsymbol{\Sigma}}_n^{-1}$$

$$\hat{\boldsymbol{\theta}}_{OLS}(\boldsymbol{X},\boldsymbol{y}) = \boldsymbol{\theta}_{\star} + \frac{1}{n}\hat{\boldsymbol{\Sigma}}_{n}^{-1}\boldsymbol{X}^{\mathsf{T}}\boldsymbol{\varepsilon}$$
"signal" "noise"

In particular:

• Bias:
$$\mathbb{E}_{\pmb{\varepsilon}} \left[\hat{\pmb{\theta}}_{OLS}(\pmb{X}, \pmb{y}) \right] = \pmb{\theta}_{\star}$$
 "Unbiased"

• Variance:
$$\operatorname{Var}_{\boldsymbol{\varepsilon}} \left[\hat{\boldsymbol{\theta}}_{OLS}(\boldsymbol{X}, \boldsymbol{y}) \right] = \frac{\sigma^2}{n} \hat{\boldsymbol{\Sigma}}_n^{-1}$$

Hence, informally:

$$\hat{\boldsymbol{\theta}}_{OLS} o \boldsymbol{ heta}_{\star}$$
 as $n o \infty$ "Consistency"

We now look at the population risk.

We now look at the population risk. First, note that for fixed $\theta \in \mathbb{R}^d$:

$$\mathcal{R}(\boldsymbol{\theta}) = \mathbb{E}_{\mathbf{y}} \left[\frac{1}{n} ||\mathbf{y} - \mathbf{X}\boldsymbol{\theta}||_{2}^{2} \right]$$

We now look at the population risk. First, note that for fixed $\theta \in \mathbb{R}^d$:

$$\mathcal{R}(\boldsymbol{\theta}) = \mathbb{E}_{\boldsymbol{y}} \left[\frac{1}{n} ||\boldsymbol{y} - \boldsymbol{X}\boldsymbol{\theta}||_{2}^{2} \right] = \mathbb{E}_{\boldsymbol{\varepsilon}} \left[\frac{1}{n} ||\boldsymbol{X}\boldsymbol{\theta}_{\star} + \boldsymbol{\varepsilon} - \boldsymbol{X}\boldsymbol{\theta}||_{2}^{2} \right]$$

We now look at the population risk. First, note that for fixed $\theta \in \mathbb{R}^d$:

$$\mathcal{R}(\boldsymbol{\theta}) = \mathbb{E}_{\mathbf{y}} \left[\frac{1}{n} ||\mathbf{y} - \mathbf{X}\boldsymbol{\theta}||_{2}^{2} \right] = \mathbb{E}_{\boldsymbol{\varepsilon}} \left[\frac{1}{n} ||\mathbf{X}\boldsymbol{\theta}_{\star} + \boldsymbol{\varepsilon} - \mathbf{X}\boldsymbol{\theta}||_{2}^{2} \right]$$
$$= (\boldsymbol{\theta}_{\star} - \boldsymbol{\theta})^{\mathsf{T}} \hat{\boldsymbol{\Sigma}}_{n} (\boldsymbol{\theta}_{\star} - \boldsymbol{\theta}) + \sigma^{2}$$

We now look at the population risk. First, note that for fixed $\theta \in \mathbb{R}^d$:

$$\mathcal{R}(\boldsymbol{\theta}) = \mathbb{E}_{\mathbf{y}} \left[\frac{1}{n} ||\mathbf{y} - \mathbf{X}\boldsymbol{\theta}||_{2}^{2} \right] = \mathbb{E}_{\boldsymbol{\varepsilon}} \left[\frac{1}{n} ||\mathbf{X}\boldsymbol{\theta}_{\star} + \boldsymbol{\varepsilon} - \mathbf{X}\boldsymbol{\theta}||_{2}^{2} \right]$$
$$= (\boldsymbol{\theta}_{\star} - \boldsymbol{\theta})^{\mathsf{T}} \hat{\boldsymbol{\Sigma}}_{n} (\boldsymbol{\theta}_{\star} - \boldsymbol{\theta}) + \sigma^{2}$$



In the fixed design setting, $X \in \mathbb{R}^{n \times d}$ is the same for training and testing.

We now look at the population risk. First, note that for fixed $\theta \in \mathbb{R}^d$:

$$\mathcal{R}(\boldsymbol{\theta}) = \mathbb{E}_{\mathbf{y}} \left[\frac{1}{n} ||\mathbf{y} - \mathbf{X}\boldsymbol{\theta}||_{2}^{2} \right] = \mathbb{E}_{\boldsymbol{\varepsilon}} \left[\frac{1}{n} ||\mathbf{X}\boldsymbol{\theta}_{\star} + \boldsymbol{\varepsilon} - \mathbf{X}\boldsymbol{\theta}||_{2}^{2} \right]$$
$$= (\boldsymbol{\theta}_{\star} - \boldsymbol{\theta})^{\mathsf{T}} \hat{\boldsymbol{\Sigma}}_{n} (\boldsymbol{\theta}_{\star} - \boldsymbol{\theta}) + \sigma^{2}$$



In the fixed design setting, $X \in \mathbb{R}^{n \times d}$ is the same for training and testing.

Therefore, for the OLS
$$\hat{\boldsymbol{\theta}}_{OLS}(\boldsymbol{X}, \boldsymbol{y}) = \boldsymbol{\theta}_{\star} + \frac{1}{n}\hat{\boldsymbol{\Sigma}}_{n}^{-1}\boldsymbol{X}^{\top}\boldsymbol{\varepsilon}$$
:

$$\mathcal{R}(\hat{\boldsymbol{\theta}}_{OLS}) - \sigma^2 = \frac{1}{n^2} \boldsymbol{\varepsilon}^{\mathsf{T}} \boldsymbol{X} \hat{\boldsymbol{\Sigma}}_n^{-1} \boldsymbol{X}^{\mathsf{T}} \boldsymbol{\varepsilon}$$

We now look at the population risk. First, note that for fixed $\theta \in \mathbb{R}^d$:

$$\mathcal{R}(\boldsymbol{\theta}) = \mathbb{E}_{\mathbf{y}} \left[\frac{1}{n} ||\mathbf{y} - \mathbf{X}\boldsymbol{\theta}||_{2}^{2} \right] = \mathbb{E}_{\boldsymbol{\varepsilon}} \left[\frac{1}{n} ||\mathbf{X}\boldsymbol{\theta}_{\star} + \boldsymbol{\varepsilon} - \mathbf{X}\boldsymbol{\theta}||_{2}^{2} \right]$$
$$= (\boldsymbol{\theta}_{\star} - \boldsymbol{\theta})^{\mathsf{T}} \hat{\boldsymbol{\Sigma}}_{n} (\boldsymbol{\theta}_{\star} - \boldsymbol{\theta}) + \sigma^{2}$$



In the fixed design setting, $X \in \mathbb{R}^{n \times d}$ is the same for training and testing.

Therefore, for the OLS
$$\hat{\boldsymbol{\theta}}_{OLS}(\boldsymbol{X}, \boldsymbol{y}) = \boldsymbol{\theta}_{\star} + \frac{1}{n}\hat{\boldsymbol{\Sigma}}_{n}^{-1}\boldsymbol{X}^{\top}\boldsymbol{\varepsilon}$$
:

$$\mathcal{R}(\hat{\boldsymbol{\theta}}_{OLS}) - \sigma^2 = \frac{1}{n^2} \boldsymbol{\varepsilon}^{\mathsf{T}} \boldsymbol{X} \hat{\boldsymbol{\Sigma}}_n^{-1} \boldsymbol{X}^{\mathsf{T}} \boldsymbol{\varepsilon}$$



This is a random variable since $\hat{m{ heta}}_{OLS}$ is random!

First, note we can rewrite:

$$\mathcal{R}(\hat{\boldsymbol{\theta}}_{OLS}) - \sigma^2 = \frac{1}{n^2} \boldsymbol{\varepsilon}^{\mathsf{T}} \boldsymbol{X} \hat{\boldsymbol{\Sigma}}_n^{-1} \boldsymbol{X}^{\mathsf{T}} \boldsymbol{\varepsilon}$$

First, note we can rewrite:

$$\mathcal{R}(\hat{\boldsymbol{\theta}}_{OLS}) - \sigma^2 = \frac{1}{n^2} \boldsymbol{\varepsilon}^{\mathsf{T}} \boldsymbol{X} \hat{\boldsymbol{\Sigma}}_n^{-1} \boldsymbol{X}^{\mathsf{T}} \boldsymbol{\varepsilon} = \frac{1}{n^2} \mathrm{Tr} \left[\boldsymbol{\varepsilon}^{\mathsf{T}} \boldsymbol{X} \hat{\boldsymbol{\Sigma}}_n^{-1} \boldsymbol{X}^{\mathsf{T}} \boldsymbol{\varepsilon} \right]$$

First, note we can rewrite:

$$\mathcal{R}(\hat{\boldsymbol{\theta}}_{OLS}) - \sigma^2 = \frac{1}{n^2} \boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \boldsymbol{\varepsilon} = \frac{1}{n^2} \operatorname{Tr} \left[\boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \boldsymbol{\varepsilon} \right]$$
$$= \frac{1}{n^2} \operatorname{Tr} \left[\boldsymbol{\varepsilon} \boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \right]$$

First, note we can rewrite:

$$\mathcal{R}(\hat{\boldsymbol{\theta}}_{OLS}) - \sigma^2 = \frac{1}{n^2} \boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \boldsymbol{\varepsilon} = \frac{1}{n^2} \text{Tr} \left[\boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \boldsymbol{\varepsilon} \right]$$
$$= \frac{1}{n^2} \text{Tr} \left[\boldsymbol{\varepsilon} \boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \right]$$

First, note we can rewrite:

$$\mathcal{R}(\hat{\boldsymbol{\theta}}_{OLS}) - \sigma^2 = \frac{1}{n^2} \boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \boldsymbol{\varepsilon} = \frac{1}{n^2} \text{Tr} \left[\boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \boldsymbol{\varepsilon} \right]$$
$$= \frac{1}{n^2} \text{Tr} \left[\boldsymbol{\varepsilon} \boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \right]$$

$$\mathbb{E}_{\boldsymbol{\varepsilon}} \left[\mathcal{R}(\hat{\boldsymbol{\theta}}_{OLS}) \right] - \sigma^2 = \frac{1}{n^2} \operatorname{Tr} \left[\mathbb{E}_{\boldsymbol{\varepsilon}} \left[\boldsymbol{\varepsilon} \boldsymbol{\varepsilon}^{\top} \right] \boldsymbol{X} \hat{\boldsymbol{\Sigma}}_n^{-1} \boldsymbol{X}^{\top} \right]$$

First, note we can rewrite:

$$\mathcal{R}(\hat{\boldsymbol{\theta}}_{OLS}) - \sigma^2 = \frac{1}{n^2} \boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \boldsymbol{\varepsilon} = \frac{1}{n^2} \operatorname{Tr} \left[\boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \boldsymbol{\varepsilon} \right]$$
$$= \frac{1}{n^2} \operatorname{Tr} \left[\boldsymbol{\varepsilon} \boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \right]$$

$$\mathbb{E}_{\boldsymbol{\varepsilon}} \left[\mathcal{R}(\hat{\boldsymbol{\theta}}_{OLS}) \right] - \sigma^2 = \frac{1}{n^2} \operatorname{Tr} \left[\mathbb{E}_{\boldsymbol{\varepsilon}} \left[\boldsymbol{\varepsilon} \boldsymbol{\varepsilon}^{\top} \right] \boldsymbol{X} \hat{\boldsymbol{\Sigma}}_n^{-1} \boldsymbol{X}^{\top} \right]$$
$$= \frac{\sigma^2}{n^2} \operatorname{Tr} \left[\boldsymbol{X} \hat{\boldsymbol{\Sigma}}_n^{-1} \boldsymbol{X}^{\top} \right]$$

First, note we can rewrite:

$$\mathcal{R}(\hat{\boldsymbol{\theta}}_{OLS}) - \sigma^2 = \frac{1}{n^2} \boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \boldsymbol{\varepsilon} = \frac{1}{n^2} \text{Tr} \left[\boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \boldsymbol{\varepsilon} \right]$$
$$= \frac{1}{n^2} \text{Tr} \left[\boldsymbol{\varepsilon} \boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \right]$$

$$\mathbb{E}_{\boldsymbol{\varepsilon}} \left[\mathcal{R}(\hat{\boldsymbol{\theta}}_{OLS}) \right] - \sigma^2 = \frac{1}{n^2} \operatorname{Tr} \left[\mathbb{E}_{\boldsymbol{\varepsilon}} \left[\boldsymbol{\varepsilon} \boldsymbol{\varepsilon}^{\top} \right] \boldsymbol{X} \hat{\boldsymbol{\Sigma}}_n^{-1} \boldsymbol{X}^{\top} \right]$$
$$= \frac{\sigma^2}{n^2} \operatorname{Tr} \left[\boldsymbol{X} \hat{\boldsymbol{\Sigma}}_n^{-1} \boldsymbol{X}^{\top} \right] = \frac{\sigma^2}{n^2} \operatorname{Tr} \left[\boldsymbol{X}^{\top} \boldsymbol{X} \hat{\boldsymbol{\Sigma}}_n^{-1} \right]$$

First, note we can rewrite:

$$\mathcal{R}(\hat{\boldsymbol{\theta}}_{OLS}) - \sigma^2 = \frac{1}{n^2} \boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \boldsymbol{\varepsilon} = \frac{1}{n^2} \text{Tr} \left[\boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \boldsymbol{\varepsilon} \right]$$
$$= \frac{1}{n^2} \text{Tr} \left[\boldsymbol{\varepsilon} \boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \right]$$

$$\mathbb{E}_{\boldsymbol{\varepsilon}} \left[\mathcal{R}(\hat{\boldsymbol{\theta}}_{OLS}) \right] - \sigma^2 = \frac{1}{n^2} \operatorname{Tr} \left[\mathbb{E}_{\boldsymbol{\varepsilon}} \left[\boldsymbol{\varepsilon} \boldsymbol{\varepsilon}^{\top} \right] \boldsymbol{X} \hat{\boldsymbol{\Sigma}}_n^{-1} \boldsymbol{X}^{\top} \right]$$

$$= \frac{\sigma^2}{n^2} \operatorname{Tr} \left[\boldsymbol{X} \hat{\boldsymbol{\Sigma}}_n^{-1} \boldsymbol{X}^{\top} \right] = \frac{\sigma^2}{n^2} \operatorname{Tr} \left[\boldsymbol{X}^{\top} \boldsymbol{X} \hat{\boldsymbol{\Sigma}}_n^{-1} \right]$$

$$= \frac{\sigma^2}{n} \operatorname{Tr} [I_d]$$

First, note we can rewrite:

$$\mathcal{R}(\hat{\boldsymbol{\theta}}_{OLS}) - \sigma^2 = \frac{1}{n^2} \boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \boldsymbol{\varepsilon} = \frac{1}{n^2} \text{Tr} \left[\boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \boldsymbol{\varepsilon} \right]$$
$$= \frac{1}{n^2} \text{Tr} \left[\boldsymbol{\varepsilon} \boldsymbol{\varepsilon}^{\top} X \hat{\boldsymbol{\Sigma}}_n^{-1} X^{\top} \right]$$

$$\mathbb{E}_{\boldsymbol{\varepsilon}} \left[\mathcal{R}(\hat{\boldsymbol{\theta}}_{OLS}) \right] - \sigma^2 = \frac{1}{n^2} \operatorname{Tr} \left[\mathbb{E}_{\boldsymbol{\varepsilon}} \left[\boldsymbol{\varepsilon} \boldsymbol{\varepsilon}^{\top} \right] \boldsymbol{X} \hat{\boldsymbol{\Sigma}}_n^{-1} \boldsymbol{X}^{\top} \right]$$

$$= \frac{\sigma^2}{n^2} \operatorname{Tr} \left[\boldsymbol{X} \hat{\boldsymbol{\Sigma}}_n^{-1} \boldsymbol{X}^{\top} \right] = \frac{\sigma^2}{n^2} \operatorname{Tr} \left[\boldsymbol{X}^{\top} \boldsymbol{X} \hat{\boldsymbol{\Sigma}}_n^{-1} \right]$$

$$= \frac{\sigma^2}{n} \operatorname{Tr} [I_d] = \sigma^2 \frac{d}{n}$$

Therefore, we have the following final result for the excess risk of OLS

$$\mathbb{E}_{\boldsymbol{\varepsilon}} \left[\mathcal{R}(\hat{\boldsymbol{\theta}}_{OLS}) \right] - \sigma^2 = \sigma^2 \frac{d}{n}$$

Therefore, we have the following final result for the excess risk of OLS

$$\mathbb{E}_{\boldsymbol{\varepsilon}} \left[\mathcal{R}(\hat{\boldsymbol{\theta}}_{OLS}) \right] - \sigma^2 = \sigma^2 \frac{d}{n}$$

Remarks:

- Excess risk is proportional to the noise level $\mathbb{E}[\varepsilon^2] = \sigma^2$.
- Excess risk is proportional to the data dimension.
- To achieve excess risk $\Delta \mathcal{R} < \delta$, need:

$$n > \frac{\sigma^2 d}{\delta}$$

samples.